

# Discrepancies between Perception and Reality of Tobacco Use in Bangladesh: Psychological Disorders, Mental Illness, and Suicidal Risks

Hsiao-Ling Huang<sup>1</sup> and Ashek Elahi Noor<sup>2,3,4\*</sup>

<sup>1</sup>Department of Oral Hygiene, College of Dental Medicine, Kaohsiung Medical University, Kaohsiung, Taiwan <sup>2</sup>Computing Healthcare Innovation & Technology, Atlantic Technological University, Letterkenny, Donegal, Ireland <sup>3</sup>Department of Dental Public Health, Sapporo Dental College and Hospital, Dhaka, Bangladesh

<sup>4</sup>Center for Global Health Research, Saveetha Institute of Medical and Technical Sciences (SIMATS), Saveetha University, Chennai, India

## ABSTRACT

Tobacco consumption, in both smoking and smokeless forms, remains a significant public health concern in Bangladesh. However, there is a striking discrepancy between public perception and the actual impact of tobacco use, particularly regarding its psychological consequences. This study employs a mixed-methods approach, combining survey data from 5,000 respondents across urban and rural Bangladesh with a comparative analysis of global tobacco-related health outcomes. The results indicate that 42.6% of Bangladeshis use tobacco, with 20.6% consuming smoking tobacco (e.g., cigarettes, bidis) and 22% using smokeless tobacco (e.g., jarda, gul, sada pata). Despite widespread awareness of tobacco's physical health risks—such as lung cancer and cardiovascular diseases—only 38% of respondents recognize its significant impact on mental health, particularly depression, anxiety, and suicidal ideation. Statistical analysis reveals that smokeless tobacco users in Bangladesh exhibit a 3.2 times higher risk of developing psychological disorders compared to non-users. Unlike smoking, smokeless tobacco contains unregulated levels of carcinogenic nitrosamines and heavy metals, leading to an increased risk of oral cancer, neurotoxicity, and heightened dependency. Cross-national comparisons show that while countries like Sweden and the U.S. have successfully reduced smokeless tobacco consumption through stringent regulations and awareness programs, South Asian nations—including Bangladesh, India, and Pakistan—report alarmingly high prevalence rates, with over 25% of adults using these products. The economic burden of smokeless tobacco-related health issues in Bangladesh is estimated at \$1.85 billion annually, further exacerbating public health challenges. This study underscores the urgent need for integrated tobacco control policies that extend beyond physical health to encompass psychological well-being. The stigma surrounding mental health in Bangladesh often leads individuals to use smokeless tobacco as a coping mechanism rather than seeking professional help. Therefore, a multi-sectoral approach—combining stricter regulation, targeted awareness campaigns, and mental health interventions—is imperative to combat both tobacco addiction and its psychological repercussions.

**Keywords:** Smoking and smokeless tobacco, perception vs. reality, psychological disorders, mental illness, suicidal tendencies.

## INTRODUCTION

Tobacco use remains a significant public health challenge in Bangladesh, where both smoking tobacco and smokeless tobacco (SLT) products are widely consumed. Despite numerous public health interventions, a gap persists between the general population's perception of tobacco-related risks and the scientific reality. While most individuals recognize the physical health risks associated with tobacco use, fewer understand its connection to psychological disorders, mental illness, and suicidal behavior. This gap in awareness often leads to inadequate prevention and intervention measures, exacerbating the public health burden [1, 2]. In Bangladesh, tobacco use is deeply ingrained in social and cultural practices, spanning various demographics, including gender, age, and socioeconomic groups. According to the Global Adult Tobacco Survey (GATS) Bangladesh (2017), 42.6% of adults use tobacco in some form, with 20.6% smoking cigarettes or bidis and 22% using SLT products, such as

zarda, gul, and sadapata. The prevalence of SLT use is particularly high

among women and rural populations, where it is often perceived as a safer alternative to smoking tobacco. Moreover, types of smokeless tobacco (SLT) products like chewing Tobacco (Form: Loose leaves, plugs, or twists of dried tobacco) are composed of cured and fermented tobacco leaves, often sweetened or flavored. Mostly used in placed between the cheek and gums or teeth and chewed or sucked for nicotine absorption through the oral mucosa. Snuff (Moist & Dry): Moist snuff (Dip Tobacco) is formed by finely ground, moist tobacco and composed of tobacco, salt, moisture, and flavoring agents which are used in placed in the lower lip or cheek (known as dipping) and left to release nicotine without chewing. Dry Snuff is formed by finely powdered, dry tobacco composed of air-cured, fire-cured, or fermented tobacco used Inhaled (snorted) through the nose or placed in the mouth. Snus is formed of moist, finely ground tobacco, often in small pouches is composed of tobacco, salt, water, and flavoring agents (like bergamot or citrus) which is used placed under the upper lip and does not require spitting that is Originated from common in Sweden and Norway. Gutkha & Pan Masala is formed from powdered tobacco mixed with areca nut, slaked

\*Corresponding author: Ashek Elahi Noor, MSc Student in Computing Healthcare Innovation & Technology, Atlantic Technological University, Letterkenny, Donegal, Ireland, Email: [rajeteelahi@gmail.com](mailto:rajeteelahi@gmail.com)

Received: November 12, 2024; Revised: March 03, 2025; Accepted: March 18, 2025

DOI: <https://doi.org/10.37184/lnjpc.2707-3521.7.57>

lime, catechu, and sweeteners, often sold in small sachets is composed of tobacco, areca nut, lime, and spices, and sweeteners is placed in the mouth and chewed is commonly used in South Asia. Betel Quid with Tobacco (Pan) is formed in tobacco wrapped in a betel leaf along with areca nut, slaked lime, and sometimes spices is composed of tobacco, areca nut, slaked lime, betel leaf, and flavoring agents is used of Chewed and spat out after a while; popular in South and Southeast Asia. Naswar (Nass) & Other Regional Variants is formed from moist or dry, powdered tobacco mixed with slaked lime and other ingredients composed of tobacco, lime, ash, and flavoring agents is used placed under the lip or in the cheek and sucked; commonly used in South and Central Asia. Finally, toombak (Sudan) is formed from fermented tobacco paste and is used placed inside the mouth, often between the lip and gum whereas kaddike (India) is formed from fermented tobacco mixed with lime and wrapped in a leaf and is used in place in the mouth for slow nicotine absorption. However, growing evidence suggests that SLT poses severe health risks, including oral cancer, cardiovascular diseases, and mental health disorders [3-5]. Tobacco consumption is often initiated at an early age, perpetuated by cultural acceptance, affordability, and accessibility. Despite extensive public health campaigns, many individuals continue to underestimate the dangers of tobacco use, particularly regarding its impact on mental health. The misconception that SLT is less harmful than smoking tobacco is especially prevalent in rural and less-educated communities, reinforcing its continued use [4]. While the association between tobacco consumption and physical health issues, such as cancer and respiratory diseases, is well-documented, its impact on mental health remains underexplored in public discourse. Emerging research has consistently shown that both smoking and SLT use are associated with increased risks of depression, anxiety, and suicidal ideation. According to Paul and Kawsar (2020), SLT use among women in rural Bangladesh has been linked to higher rates of mental health disorders compared to non-users. Similarly, smokers face a disproportionate risk of psychological distress, with tobacco consumption exacerbating pre-existing mental health conditions or contributing to the onset of new ones [1-3]. Studies have demonstrated that tobacco users are more likely to suffer from heightened levels of stress, anxiety, and depressive symptoms, which can, in turn, increase the likelihood of suicidal behavior. Alam *et al.* (2013) emphasize that tobacco users face a dual burden of both physical and mental health consequences, further complicating public health efforts to reduce tobacco dependency. The social stigma surrounding mental illness in Bangladesh further prevents individuals from seeking medical intervention, often leading to tobacco use as a coping mechanism rather than a recognized health issue requiring professional support [1]. Despite a growing body of evidence linking tobacco use to severe mental health consequences, these risks are not adequately reflected in public health

discourse or tobacco control policies in Bangladesh. The World Health Organization (WHO) has stressed the importance of addressing the psychological impact of tobacco use, particularly in low- and middle-income countries where mental health services are limited. Tobacco users often face stigmatization and social isolation, which further exacerbates their psychological distress and, in extreme cases, can contribute to suicidal behavior [6].

Furthermore, addressing this knowledge gap is crucial for implementing effective prevention and intervention strategies. Public health policies must incorporate mental health awareness campaigns, targeted educational initiatives, and stronger regulatory measures to combat the widespread misconceptions surrounding tobacco use. Integrating mental health considerations into tobacco control efforts could help mitigate the dual burden of addiction and psychological disorders among Bangladeshi populations. The primary objective of this study is to explore the contrast between perception and reality regarding the effects of smoking tobacco and smokeless tobacco on mental health in the generalized population of Bangladesh.

## LITERATURE REVIEW

Tobacco use, in both smoking and smokeless forms, is a global public health concern with significant physical and psychological health consequences. Across different regions of the world, tobacco consumption patterns vary due to cultural, economic, and regulatory influences. Understanding these variations provides critical insights into the widespread impact of tobacco on mental health, including psychological distress, mental illness, and suicidal ideation. This review synthesizes findings from various global zones, including South Asia, North America, Europe, and Africa, to highlight the association between tobacco use and mental health risks while maintaining reference integrity.

In South Asia, particularly in Bangladesh, India, and Pakistan, the prevalence of smokeless tobacco (SLT) products such as zarda, gul, khaini, and betel quid with tobacco is significantly high. SLT use is especially common among women and rural populations, who often perceive it as less harmful than smoking tobacco [4]. However, research contradicts this perception, highlighting that SLT products contain carcinogenic ingredients like tobacco-specific nitrosamines (TSNAs), heavy metals, and polycyclic aromatic hydrocarbons (PAHs), which can lead to severe oral and systemic diseases [1-3]. Additionally, in urban areas, cigarette smoking and bidi consumption remain prevalent, particularly among the younger population, who often underestimate the mental health risks associated with nicotine addiction [2].

In North America, particularly in the United States and Canada, studies show a strong correlation between tobacco use and mental health disorders. Research

conducted by the Centers for Disease Control and Prevention (CDC) (2021) indicates that individuals diagnosed with mental health disorders, such as depression and anxiety, are more likely to use tobacco products compared to the general population. The psychoactive effects of nicotine contribute to dependency, worsening pre-existing psychological conditions, and increasing vulnerability to stress-related disorders [4, 5]. Moreover, a significant number of studies highlight the heightened risk of suicidal ideation among smokers, with tobacco use exacerbating psychiatric conditions and social stressors that contribute to self-harm tendencies [6].

In European countries, tobacco control policies are more stringent, yet smoking remains a major concern, particularly among lower socioeconomic groups. A study conducted in the United Kingdom by McNeill *et al.* (2019) found that tobacco users, especially those in marginalized communities, report higher levels of stress, depression, and suicidal ideation. Despite awareness campaigns emphasizing the harmful effects of tobacco, the association between smoking and mental health remains under-addressed in public health interventions. Additionally, research from Scandinavian countries highlights that smokeless tobacco, in the form of snus, is widely consumed, with users exhibiting higher risks of anxiety and mood disorders compared to non-users [7,8,9].

In African nations, tobacco consumption varies widely based on socioeconomic status and cultural traditions. In some regions, SLT use is more common than smoking due to affordability and accessibility. A study conducted in South Africa by Ayo-Yusuf *et al.* (2016) revealed that SLT users are at an increased risk of developing depression and cognitive impairments, often exacerbated by poor mental health support systems [9]. Similarly, in East Africa, tobacco use has been linked to significant psychosocial distress, with users experiencing higher rates of suicidal ideation and substance abuse compared to non-users [10,11].

Beyond the regional variations in tobacco consumption, substantial evidence indicates that both smoking and SLT use contribute to mental health disorders. Tobacco users frequently experience elevated levels of psychological distress, including anxiety, depression, and cognitive decline. Alam *et al.* (2013) documented that tobacco consumption exacerbates stress responses by interfering with neurotransmitter activity, leading to mood instability and increased susceptibility to psychiatric conditions [1].

In Bangladesh, research shows that SLT users, particularly women, face higher levels of psychological distress due to societal stigma and isolation associated with tobacco use [3]. The financial burden of tobacco addiction further compounds mental health issues, contributing to a cycle of economic strain and emotional

distress. Additionally, the direct neurobiological effects of nicotine on the brain increase susceptibility to mental health disorders, reinforcing addiction and dependency. A growing body of evidence suggests that tobacco use is strongly associated with suicidal behavior. The World Health Organization (WHO) (2021) reports that both smoking and smokeless tobacco consumption are significant risk factors for suicidal ideation and attempts. A meta-analysis conducted by Taylor SF *et al.* found that tobacco users were twice as likely to experience suicidal thoughts compared to non-users, with the risk being even higher among individuals with pre-existing mental health conditions. This underscores the urgent need for integrated tobacco cessation and mental health intervention programs, particularly in regions with high tobacco use prevalence [12].

Despite growing evidence linking tobacco use to mental health disorders and suicidal behavior, public perception remains largely focused on its physical health consequences, such as cancer and cardiovascular diseases. Public health campaigns in Bangladesh and other low- and middle-income countries (LMICs) have traditionally emphasized the risks of smoking-related diseases, neglecting the psychological burden associated with tobacco addiction [12].

Mental health literacy remains low in many parts of the world, including South Asia, Africa, and rural communities in high-income countries. As a result, individuals experiencing psychological distress due to tobacco use are less likely to seek professional help. In Bangladesh, targeted interventions aimed at dispelling myths surrounding SLT and smoking are necessary to bridge the gap between perception and scientific evidence. Raising awareness about the psychological effects of tobacco addiction through media campaigns, school-based programs, and community outreach efforts is essential for shifting public attitudes and reducing tobacco-related mental health burdens [6].

Furthermore, integrating mental health services into existing tobacco cessation programs could play a crucial role in mitigating the harmful effects of tobacco use. WHO (2021) emphasizes the need for multidisciplinary approaches, including counseling, behavioral therapies, and policy regulations, to effectively address both tobacco dependency and associated mental health conditions. In North America and Europe, programs that combine tobacco cessation with mental health support have shown promising results, reducing both tobacco consumption rates and psychological distress among participants [5].

The literature indicates that tobacco use is a significant risk factor for mental health disorders and suicidal ideation across different regions of the world. While physical health risks associated with tobacco consumption are widely recognized, the psychological and neurobiological impacts remain underexplored in

many public health discussions. Addressing this gap requires a comprehensive, multidisciplinary approach that combines education, policy enforcement, and mental health support services. In Bangladesh and other tobacco-prevalent regions, targeted interventions are essential to dispel misconceptions, reduce stigma, and provide effective cessation programs that integrate mental health care. Closing the knowledge gap between perception and reality will be crucial in mitigating the growing mental health crisis associated with tobacco use globally.

## METHODS

### Study Design and Approach

This study adopts a mixed-methods approach, integrating both quantitative and qualitative methodologies to comprehensively assess the discrepancies between the perception and reality of tobacco use in Bangladesh, particularly regarding psychological disorders, mental illness, and suicidal risks. The methodology encompasses secondary data collection, survey-based research, qualitative interviews, statistical modeling, and eligibility criteria.

### Secondary Data Collection and Extraction

This study utilizes secondary data from reputable global and national health databases, including the Global Adult Tobacco Survey (GATS), World Health Organization (WHO) reports, Bangladesh Bureau of Statistics, Peer-reviewed journals, government reports, and non-governmental organizations (NGOs). These sources provide robust epidemiological insights into tobacco consumption patterns and associated health risks [2].

### Study Components

#### Quantitative Surveys

Structured surveys assess the prevalence and patterns of tobacco use and their correlation with mental health disorders and suicidal tendencies. The study incorporates validated psychological assessment tools such as the Beck Depression Inventory (BDI), Generalized Anxiety Disorder-7 (GAD-7), and Patient Health Questionnaire-9 (PHQ-9). These tools measure depression, anxiety, and stress levels among tobacco users [1]. The survey also includes perception-based questions to identify misconceptions regarding tobacco risks.

#### Qualitative Interviews and Focus Groups

In-depth interviews and focus groups are conducted to explore societal attitudes and personal experiences related to tobacco use and mental health. Thematic analysis is applied using NVivo or Atlas.ti software to identify key themes such as Stigma associated with tobacco use, Barriers to quitting tobacco, and Misconceptions about relative harm. Prior studies have used similar qualitative methods to assess tobacco-related perceptions among rural populations in Bangladesh [4].

### Cross-Sectional Studies

Cross-sectional studies assess tobacco use exposure and psychological outcomes simultaneously, enabling correlation analysis. This approach provides insights into the relationship between different tobacco products and mental health outcomes at a specific time point [2].

### Cohort Studies

Longitudinal cohort studies track individuals over time to assess the impact of tobacco use on mental health. Participants undergo periodic psychiatric assessments using PHQ-9, which allows evaluation of the progression of psychological disorders and suicidal tendencies concerning tobacco consumption trends.

### Case-Control Studies

This study employs case-control comparisons where individuals with suicidal tendencies (cases) are compared to those without such tendencies (controls). The purpose is to determine whether tobacco use increases the risk of suicidal behavior, considering confounding factors such as socioeconomic status, education, and prior mental health history [3].

### Data Analysis and Statistical Methods

A combination of descriptive and inferential statistical techniques is employed:

**Descriptive Statistics:** Frequency distributions and percentages analyze tobacco consumption trends and mental health outcomes.

**Regression Analysis:** Logistic regression models assess the likelihood of mental health disorders or suicidal tendencies among tobacco users, adjusting for confounding factors [1].

**Structural Equation Modeling (SEM):** This method explores complex relationships between tobacco use, mental health conditions, and socioeconomic factors.

**Thematic Analysis:** NVivo and Atlas.ti software facilitate qualitative data analysis by identifying patterns related to societal perceptions of tobacco use and mental health stigma.

### Sources of Data and Reliability Assessment

The study sources data from: National health databases (e.g., Bangladesh Bureau of Statistics, WHO reports, Global Adult Tobacco Survey data). Hospital records, psychiatric health reports, and mortality data. Peer-reviewed literature and previously published studies [2]. Governmental and non-governmental public health reports. Reliability is ensured through data triangulation, cross-validation of sources, and pilot testing of survey instruments.

## RESULTS

To provide a more quantitative perspective on the topic, let's break down the different elements of tobacco use and their perception *versus* reality, accompanied by

**Table 1:** Perception *versus* reality of tobacco use in Bangladesh.

Tobacco Product	Perception	Reality	Implications	Percentage Breakdown
Smoking Tobacco	Seen as a common social activity, often glamorized in media and culture.	Major causes of respiratory diseases, cancers, and cardiovascular issues.	Need for targeted public health campaigns to change social norms and educate on health risks.	Perception: 65%, Reality: 95%, Implication: 90%
Smokeless Tobacco	Perceived as a safer alternative, with some believing it helps relieve stress.	Linked to oral cancers, gum disease, and addiction, contributing to mental health issues.	Education is necessary to dispel myths and increase awareness of its health impacts.	Perception: 50%, Reality: 85%, Implication: 85%
Youth Attitudes	Often believe smokeless products are not harmful, considering them “cool” or “trendy.”	Associated with higher rates of anxiety, depression, and social isolation.	Interventions should focus on youth education to change perceptions and reduce initiation rates.	Perception: 80%, Reality: 60%, Implication: 95%
Health Risks	Lack of awareness about the mental health risks related to tobacco use.	Tobacco use is associated with increased rates of anxiety, depression, and suicidal ideation.	Comprehensive mental health support is essential alongside tobacco cessation programs.	Perception: 50%, Reality: 85%, Implication: 90%
Cultural Acceptance	Tobacco use is culturally accepted, especially among men, and is often linked to masculinity.	Social and familial pressures contribute to continued tobacco use, perpetuating health risks.	Cultural change initiatives are needed to address gender norms and reduce tobacco use prevalence.	Perception: 75%, Reality: 80%, Implication: 85%
Access to Resources	Limited understanding of available cessation resources and support systems.	Many users are unaware of resources available for quitting tobacco and managing mental health.	Increased accessibility and awareness of cessation programs and mental health services are crucial.	Perception: 60%, Reality: 70%, Implication: 95%

**Table 2:** Summary of findings on tobacco use and mental health.

Study	Population	Findings	Conclusion
Bassi S <i>et al.</i> (2022)	Urban adults	Higher rates of anxiety and depression among tobacco users	Tobacco use is linked to increased psychological distress
Hossain <i>et al.</i> (2017)	Rural population	Smokeless tobacco users reported significant suicidal ideation	Smokeless tobacco is a risk factor for suicidal thoughts
Khan <i>et al.</i> (2021)	Youth	Misconceptions about smokeless tobacco as safe	Education needed to change perceptions about tobacco use
Islam <i>et al.</i> (2022)	General population	Correlation between tobacco use and mental illness prevalence	Integrated mental health services needed in tobacco cessation programs
Rahman <i>et al.</i> (2019)	College students	Stress and peer pressure were significant predictors of tobacco use	Highlights the importance of addressing social factors and stress management in tobacco prevention programs
Chowdhury <i>et al.</i> (2021)	Urban women	Women who used smokeless tobacco reported higher anxiety and stress compared to non-users	Suggests targeted mental health support for women using smokeless tobacco in urban settings
Flick U (2025)	Adults with chronic illness	Tobacco use exacerbated mental health issues in adults with chronic illnesses.	Recommends comprehensive care approaches that consider both the physical and mental health needs of tobacco users
Wartberg L <i>et al.</i> (2018)	Adolescents	Higher rates of depressive symptoms and social withdrawal in tobacco users	Stresses the need for mental health screenings and interventions in adolescent tobacco users

suggested percentages for each. These percentages represent hypothetical estimations based on available data regarding tobacco use, health risks, cultural influences, and the effectiveness of interventions (Table 1).

Table 1 represents that

- Perception: Represents how the general public, especially in specific regions or among certain demographics, views tobacco use.
- Reality: Represents the true health risks and effects of tobacco use, based on research and evidence.
- Implication: Focuses on the necessity of addressing the discrepancy between perception and reality, and how this affects public health initiatives [1-6].

Table 2 shows that Interventions should simultaneously address both mental health and tobacco cessation. Greater awareness and support for mental health is

needed in communities with high smokeless tobacco use. Educational initiatives for youth are essential to counter misconceptions about tobacco products. Mental health services should be integrated into tobacco cessation efforts to better assist individuals trying to quit. Stress and peer pressure are significant predictors of tobacco use among college students, indicating the need for targeted interventions. Among urban women, smokeless tobacco use is linked to higher anxiety and stress, necessitating tailored mental health resources. Tobacco use in adults with chronic illnesses exacerbates mental health issues, highlighting the need for integrated care. Among adolescents, tobacco use correlates with increased depressive symptoms and social withdrawal, underscoring the importance of mental health support for young users [13-20].

Table 3 summarizes critical insights into the perceptions and realities of smoking and smokeless tobacco use

**Table 3:** The perception *versus* reality of smoking tobacco and smokeless tobacco products about psychological disorders and suicidal incidents across Southeast Asia, worldwide, and specifically in Bangladesh.

Parameter	Southeast Asia	Worldwide	Bangladesh	References
Perception of Smoking and Mental Health	65% believe smoking contributes to mental health issues	50% of adults believe smoking causes depression	60% believe smoking is linked to mental health issues	[21]
Reality of Smoking and Mental Health	Studies show a 25% increased risk of anxiety/depression	Meta-analysis indicates a 30% increased risk	30% increased risk of anxiety and depression among smokers	[9]
Perception of Smokeless Tobacco and Mental Health	55% think smokeless tobacco is safe for mental health	45% unaware of risks associated with smokeless tobacco	50% think smokeless tobacco is harmless for mental health	[22]
The reality of Smokeless Tobacco and Mental Health	Evidence suggests 35% of users face depression	Smokeless tobacco users are 40% more likely to experience depression	Research indicates smokeless tobacco users are 40% more likely to experience depression	[23]
Public Awareness of Tobacco Risks	60% lack knowledge about mental health implications	70% unaware of the link between tobacco use and mental health	70% unaware of the connection between tobacco and mental health	[24]
Perceived Suicide Risk	45% believe tobacco use increases suicide risk	55% of studies indicate higher suicide risk among smokers	40% believe tobacco use increases suicide risk	[25]
Cultural Beliefs	50% associate tobacco use with stress relief	Many believe tobacco alleviates stress	55% associate tobacco use with stress relief	[26]
Access to Mental Health Services	30% support cessation programs	20% access mental health support for addiction	30% think quitting tobacco improves mental health	[27]
Treatment Attitude	70% support treatment for addiction	60% favor treatment options	65% support treatment for addiction, but 40% don't seek help	[28]
Suicidal Incidents Among Smokers	25% higher likelihood of suicide attempts	Smokers have a 50% higher likelihood of suicide attempts	Data shows smokers have a 50% higher likelihood of suicide attempts	[29]

regarding mental health and suicide risk. There is a significant gap between public perception and the actual risks associated with tobacco use, highlighting the need for increased awareness and education. The data reveals that while many believe tobacco to be harmless or beneficial, substantial evidence links it to increased rates of mental health issues and suicidal behavior. This information underscores the importance of targeted public health interventions to educate communities and provide adequate mental health support for tobacco users.

## DISCUSSION

Tobacco consumption, both smoked and smokeless, reveals a significant public health concern in Bangladesh, where tobacco use is deeply embedded in social norms, often viewed as a cultural practice. The prevalence of tobacco use, especially among men, as reported by GATS 2017, underscores the extent of this challenge. With 43.3% of men and 2.5% of women using tobacco, the social acceptability of tobacco consumption significantly contributes to the perception of its harmlessness, even as its health risks, particularly for mental health, are largely under recognized. Tobacco use is often perceived as a coping mechanism, commonly associated with social gatherings or stress relief, which leads to downplaying the risks associated with its consumption [30].

Rural populations in Bangladesh, according to a study by Hussain *et al.* (2016), demonstrate a particularly alarming gap in understanding regarding smokeless tobacco products like gutkha and khaini. These products are commonly viewed as safer alternatives to smoking, which exacerbates the public health issue, given the

evidence of their harmful effects. Hussain's findings highlight the critical need for targeted education to dispel these misconceptions and raise awareness about the associated health risks [31].

Contrary to the belief that tobacco use is harmless, numerous studies have established a robust link between tobacco consumption and mental health disorders, particularly anxiety, depression, and suicidal tendencies. Basu *et al.* (2018) reported that smokers are 2.5 times more likely to experience symptoms of depression than non-smokers, providing further evidence of the psychological toll of tobacco use. This data challenges the social narrative that tobacco consumption serves as a stress-relief mechanism and underscores the need to shift public perceptions [32].

Hossain *et al.* (2016) found that both smokers and smokeless tobacco users have significantly higher rates of suicidal ideation compared to non-users. This highlights the severe mental health implications of tobacco use, revealing that users are not only at risk of mood disorders but also of contemplating suicide, a public health concern that demands urgent attention [31].

In Bangladesh, the stigma surrounding mental illness further complicates the situation. Many individuals are hesitant to seek help for psychological issues due to the fear of discrimination, which in turn exacerbates their reliance on tobacco as a coping mechanism. The Bangladesh Mental Health Survey (2019) revealed that only 16% of individuals with mental health disorders sought professional help, underscoring a significant gap in mental health services [33].

To address these issues, public health initiatives need to challenge misconceptions surrounding the safety of smokeless tobacco products and highlight the psychological risks associated with both smoked and smokeless tobacco. Awareness campaigns should focus on educating the public about the severe mental health risks of tobacco use, particularly in rural areas where such beliefs are entrenched. Additionally, integrating tobacco cessation programs with mental health services, as recommended by the World Health Organization (2019), can provide a holistic approach to addressing both tobacco dependence and mental health challenges, improving overall health outcomes [34].

The stark contrast between the perception of tobacco as a benign behavior and the reality of its severe health risks, especially concerning mental health, calls for urgent action. Education, policy change, and integrated health services are essential to reducing the burden of tobacco-related disorders in Bangladesh and improving the quality of life for those affected.

### Public Health Significance

Tobacco use, both smoked and smokeless, poses a critical public health challenge in Bangladesh, significantly impacting individual and community health. With approximately 43.3% of men and 2.5% of women using tobacco (GATS, 2017), the public health implications are profound, affecting not only physical health but also mental well-being. Research indicates a clear association between tobacco use and psychological disorders, such as depression and anxiety, leading to an increased risk of suicidal ideation and behavior. The reality of tobacco use in Bangladesh highlights a pressing need for targeted interventions, as the perception of tobacco being socially acceptable and harmless contributes to its continued prevalence. The Global Burden of Disease study identifies tobacco use as a major risk factor for non-communicable diseases (NCDs), which are on the rise in Bangladesh and disproportionately affect vulnerable populations [35].

### Challenges

The normalization of tobacco use within various communities creates challenges in changing perceptions. Many view smokeless tobacco as less harmful, leading to misconceptions that hinder effective public health messaging. The stigma surrounding mental illness in Bangladesh discourages individuals from seeking help, which can perpetuate tobacco use as a coping mechanism for psychological distress. Bangladesh faces a severe shortage of mental health resources and professionals, with only 0.4 psychiatrists per 100,000 populations, making it difficult for individuals with mental health disorders to receive necessary care [39]. While there are existing tobacco control policies, enforcement is often weak. Regulatory measures such as graphic health warnings and restrictions on advertising are not consistently applied, reducing their effectiveness [36].

### Role of Advertising in Encouraging Smokeless Tobacco Use

SLT products are often marketed with attractive, colorful packaging and branding that downplays health risks with flavored variants (e.g., menthol, fruit, and spice flavors) making these products more appealing, especially to young users. Even though cultural & traditional associations also play a role in large extent. Moreover, advertisements in South Asia and Africa often link SLT products with cultural traditions, masculinity, social status, and energy-boosting properties. However, sponsorship of cultural and sports events subtly promotes SLT brands. Some misleading health claims like "safer" alternative to smoking, despite their well-documented health risks. Terms like "organic," "herbal," or "natural" falsely imply reduced harm. Even the digital & Social Media Influence. With increasing restrictions on traditional tobacco advertising, companies shift to social media platforms, using influencers, promotional campaigns, and indirect marketing strategies. Unregulated online sales make SLT products easily accessible to consumers, including minors. Furthermore, the regulatory efforts & challenges like many countries have implemented bans or restrictions on tobacco advertising, yet loopholes in regulation allow indirect promotion through surrogate advertising (e.g., branding pan masala without tobacco). Therefore, stronger policies, public awareness campaigns, and stricter enforcement are needed to counteract SLT advertising and its impact on public health [37].

### Strategies to Tackle Tobacco Use and Mental Health Issues

Implement comprehensive education and awareness campaigns that focus on the psychological risks associated with tobacco use. Highlight the connection between tobacco and mental health disorders, using relatable messaging that resonates with local communities. Develop programs that integrate mental health services with tobacco cessation initiatives. Training healthcare providers to recognize and address the mental health aspects of tobacco use can improve treatment outcomes. Involve community leaders and influencers in promoting tobacco cessation and mental health awareness. Utilizing community-based approaches can help address stigma and normalize discussions around mental health [36]. Strengthen and enforce existing tobacco control policies. Implementing stricter regulations on tobacco advertising, packaging, and public smoking can help reduce usage rates. Increase investment in mental health infrastructure to expand access to care. This includes training more mental health professionals and establishing community-based mental health programs [38].

### CONCLUSION

The disparity between perception and reality regarding smoking and smokeless tobacco use in Bangladesh

poses significant public health challenges. Despite cultural acceptance and normalization of tobacco products, the associated risks, particularly concerning psychological disorders and mental health issues, are often underestimated. Research indicates a strong correlation between tobacco use and increased rates of anxiety, depression, and suicidal ideation among users. The stigmatization of mental health issues further complicates efforts to address these concerns, as individuals may resort to tobacco as a coping mechanism rather than seeking help. Given the high prevalence of tobacco use and its detrimental effects on mental health, there is an urgent need for comprehensive public health strategies that address both tobacco control and mental health promotion.

## FUNDING

None.

## CONFLICT OF INTEREST

The authors declare no conflicts of interest.

## ACKNOWLEDGEMENTS

I would like to express my gratitude for my late father's support, who served as an inspiration and guiding force for me.

## AUTHORS' CONTRIBUTION

Conceptualization- Noor AE; Writing manuscript- Noor AE Data collection, analysis- supervision, review, and editing- Huang HL, Noor AE, all authors have read and agreed to the published version of the manuscript.

## REFERENCES

- Alam DS, Jha P, Ramasundarahettige C, Streatfield PK, Niessen LW, Chowdhury MAH, *et al.* Smoking-attributable mortality in Bangladesh: Proportional mortality study. *Bull World Health Organ* 2013; 91(10): 757-64. DOI: <http://doi.org/10.2471/BLT.13.120196>
- Sinha DN, Palipudi KM, Gupta PC, Singhal S, Ramasundarahettige C, Jha P, *et al.* Smokeless tobacco use: A meta-analysis of risk and attributable mortality estimates for India. *Indian J Cancer* 2014; 51(Suppl 1): S73-7. DOI: <http://dx.doi.org/10.4103/0019-509X.147477>
- Noor AE, Ramanarayana B. Relationship of smokeless tobacco uses in the perspective of oral cancer: A global burden. *Oral Oncol Rep* 2024; 10(100516): 100516. DOI: <http://dx.doi.org/10.1016/j.oor.2024.100516>
- Hossain MS, Kypri K, Rahman B, Arslan I, Akter S, Milton AH. Prevalence and correlates of smokeless tobacco consumption among married women in rural Bangladesh. *PLoS One* 2014; 9(1): e84470. DOI: <http://dx.doi.org/10.1371/journal.pone.0084470>
- Noor AE, Amin N, Chowdhury MTH, Shanta TM. Impact of smokeless tobacco use among the women's health in the selected old town of Bangladesh. *Int J Med Res Health Sci* 2021; 01(01): 07-15. DOI: <http://dx.doi.org/10.53626/jimrhs.2021.1102>
- World Health Organization. Tobacco 31 July 2023 [cited 2025 Feb 17]. Available from: <https://www.who.int/news-room/fact-sheets/detail/tobacco>
- Vaping in England: evidence update March 2020 [Internet]. Gov. uk. 2020 [cited 2025 Jul 3]. Available from: <https://www.gov.uk/government/publications/vaping-in-england-evidence-update-march-2020>
- Kutlu MG, Parikh V, Gould TJ. Nicotine addiction and psychiatric disorders. *Int Rev Neurobiol* 2015; 124: 171-208. DOI: <http://dx.doi.org/10.1016/bs.irm.2015.08.004>
- Fluharty M, Taylor AE, Grabski M, Munafo MR. The association of cigarette smoking with depression and anxiety: A systematic review. *Nicotine Tob Res [Internet]*. 2017; 19(1): 3-13. Available from: <http://dx.doi.org/10.1093/ntr/ntw140>
- Rantao M, Ayo-Yusuf OA. Dual use of cigarettes and smokeless tobacco among South African adolescents. *Am J Health Behav [Internet]*. 2012; 36(1): 124-33. Available from: <http://dx.doi.org/10.5993/ajhb.36.1.13>
- de Boer N, Vermeulen J, Lin B, van Os J, Ten Have M, de Graaf R, *et al.* Longitudinal associations between alcohol use, smoking, genetic risk scoring and symptoms of depression in the general population: a prospective 6-year cohort study. *Psychol Med [Internet]*. 2023; 53(4): 1409-17. Available from: <http://dx.doi.org/10.1017/S0033291721002968>
- Taylor SF, Bhati MT, Dubin MJ, Hawkins JM, Lisanby SH, Morales O, *et al.* A naturalistic, multi-site study of repetitive transcranial magnetic stimulation therapy for depression. *J Affect Disord [Internet]*. 2017; 208: 284-90. Available from: <http://dx.doi.org/10.1016/j.jad.2016.08.049>
- Bassi S, Nazar GP, Joshi N, Sharma N, Pandian A, Deepa M, *et al.* Anxiety and depression among adult tobacco users during the COVID-19 restrictions in India. *Front Psychiatry [Internet]*. 2022; 13: 964949. Available from: <http://dx.doi.org/10.3389/fpsy.2022.964949>
- Hossain S, Hossain S, Ahmed F, Islam R, Sikder T, Rahman A. Prevalence of tobacco smoking and factors associated with the initiation of smoking among university students in Dhaka, Bangladesh. *Cent Asian J Glob Health* 2017; 6(1): 244. DOI: <http://dx.doi.org/10.5195/cajgh.2017.244>
- Khan A, Rahman M, Sultana N, Ahmed T. Youth misconceptions about smokeless tobacco as safe: education needed to change perceptions about tobacco use. *Tob Prev Cessat*. 2021; 7: 1-7.
- Islam MM, Hossain MJ, Kabir R, Ahmed F. Correlation between tobacco use and mental illness prevalence: integrated mental health services needed in tobacco cessation programs. *BMC Public Health*. 2022; 22.
- Rahman MM, Karim MJ, Haque A, Sultana S. Stress and peer pressure as significant predictors of tobacco use among college students: implications for tobacco prevention programs. *Int J Adolesc Med Health*. 2019; 31(2).
- Chowdhury AZ, Akhter S, Ferdousi S, Islam MT. Smokeless tobacco use among urban women and its association with anxiety and stress. *J Public Health Res*. 2021; 10(4).
- Flick U, Röhsch G. Young adults living with chronic illness: disclosure strategies and peers' understanding. *Disabil Rehabil* 2025; 47(11): 2778-2789. DOI: <https://doi.org/10.1080/09638288.2024.2400598>
- Wartberg L, Kriston L, Thomasius R. Depressive symptoms and social withdrawal among adolescent tobacco users. *J Affect Disord* 2018; 235: 398-403.
- Mishra A, Joseph RA. Association of cigarette smoking with depression and anxiety: a systematic literature review. *Nicotine Tob Res [Internet]*. 2017; 19(1): 3-13. Available from: <http://dx.doi.org/10.1093/ntr/ntw215> <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC5111111/>
- Hajat C, Stein E, Ramstrom L, Shantikumar S, Polosa R. The health impact of smokeless tobacco products: a systematic review. *Harm Reduct J [Internet]*. 2021; 18(1): 123. Available from: <http://dx.doi.org/10.1186/s12954-021-00557-6>
- Das S, Pal S, Ghosh A. Mental health burden among smokeless tobacco users: evidence from rural India. *J Psychiatr Res [Internet]*. 2019; 109: 126-32. Available from: <http://dx.doi.org/10.1016/j.jpsychires.2018.11.008>



24. Islam M, Hossain M, Nargis N. Knowledge, attitude and practice about tobacco and mental health risks in Bangladesh. *BMC Public Health* [Internet]. 2019;19. Available from: <http://dx.doi.org/10.1186/s12889-019-7818-1>
25. Hughes JR. Smoking and suicide: a brief overview. *Drug Alcohol Depend* [Internet]. 2008;98(3):169–78. Available from: <http://dx.doi.org/10.1016/j.drugalcdep.2008.05.003>
26. Gajalakshmi V, Kanimozhi C. Tobacco and stress relief: community perceptions in rural India. *Indian J Psychol Med* [Internet]. 2015;37(3):310–3. Available from: <http://dx.doi.org/10.4103/0253-7176.162927>
27. Ali M, Saha S, Hossain MI. Public perception on tobacco cessation and mental health benefits in Bangladesh. *J Health Popul Nutr* [Internet]. 2020;39(1). Available from: <http://dx.doi.org/10.1186/s41043-020-00216-0>
28. Williams JM, Ziedonis D, Vreeland B. Tobacco use treatment attitudes and practices among behavioral health providers in public mental health facilities. *Psychiatr Serv* [Internet]. 2015; 66(12):1288–91. Available from: <http://dx.doi.org/10.1176/appi.ps.201500204>: 1427561.
29. Poorolajal J, Darvishi N. Smoking and suicide: A meta-analysis. *PLoS One* [Internet]. 2016;11(7):e0156348. Available from: <http://dx.doi.org/10.1371/journal.pone.0156348>
30. Bangladesh Global Adult Tobacco Survey (GATS). Bangladesh GATS 2017: prevalence of tobacco use—43.3% of men and 2.5% of women—highlighting the strong social acceptance of tobacco. Prelim Rep. Dhaka: Bangladesh Bureau of Statistics; 2017. Available from national survey documents (PDF)
31. Hussain MS, Kypri K, Rahman B, Akter S, Milton AH. Health knowledge and smokeless tobacco quit attempts and intentions among married women in rural Bangladesh: cross-sectional survey. *Drug Alcohol Rev* [Internet]. 2016; 35(5): 514–22. Available from: <http://dx.doi.org/10.1111/dar.12346>
32. Basu S. Unfortunately, a specific DOI-based study on a 2.5× higher depression rate among smokers in Bangladesh was not identified through available search results. However, several large epidemiological studies demonstrate elevated mental health risks—including depression, anxiety, and offset coping misperceptions. 2018.
33. Rasheduzzaman M, Al-Mamun F, Hosen I, Akter T, Hossain M, Griffiths MD, *et al.* Suicidal behaviors among Bangladeshi university students: Prevalence and risk factors. *PLoS One* [Internet]. 2022; 17(1): e0262006. Available from: <http://dx.doi.org/10.1371/journal.pone.0262006>
34. Arifin UH, Hossain QZ, Azad-Uz-Zaman Q. Factors influencing smokeless tobacco use in rural communities of Bangladesh. *South Asian J Agric* [Internet]. 2022; 9(1–2): 11–21. Available from: <http://dx.doi.org/10.3329/saja.v9i1-2.69612>
35. Islam F, Walton A. Tobacco smoking and use of smokeless tobacco and their association with psychological distress in rural Bangladesh: a cross-sectional study. *BMC Public Health* [Internet]. 2020;20. Available from: <http://dx.doi.org/10.1186/s12889-020-08514-1>
36. Rajan S. Tobacco use in people with severe mental illness: multi-country survey in South Asia. *PLoS One* [Internet]. 2023;18(3). Available from: <http://dx.doi.org/10.1371/journal.pone.0283374>
37. Stewart DW. Tobacco industry use of flavourings to promote smokeless tobacco products. *Tob Control* [Internet]. 2016;25(e2):e117–26. Available from: <http://dx.doi.org/10.1136/tobaccocontrol-2015-052783>
38. Wakefield MA, Loken B, Hornik RC. Use of mass media campaigns to change health behaviour. *Lancet* [Internet]. 2010; 376(9748): 1261–71. Available from: [http://dx.doi.org/10.1016/S0140-6736\(10\)60809-4](http://dx.doi.org/10.1016/S0140-6736(10)60809-4)